



TITLE:

14.  $\beta$  ORDERING PROCESS IN A  
QUENCHED TRICRITICAL  
SYSTEM(基研研究会「パターン形成  
,その運動と統計」,研究会報告)

AUTHOR(S):

Ohta, T.; Kawasaki, K.; Sato, A.; Enomoto, Y.

---

CITATION:

Ohta, T. ...[et al]. 14.  $\beta$  ORDERING PROCESS IN A QUENCHED TRICRITICAL SYSTEM(基研研究会「パターン形成,その運動と統計」,研究会報告). 物性研究 1987, 49(1): 43-43

ISSUE DATE:

1987-10-20

URL:

<http://hdl.handle.net/2433/92851>

RIGHT:

## 14. /3 ORDERING PROCESS IN A QUENCHED TRICRITICAL SYSTEM

T. Ohta, K. Kawasaki, A. Sato

Department of Physics, Kyushu University, Fukuoka 812, Japan

and Y. Enomoto

Department of Applied Physics, Nagoya University Nagoya, 464 Japan

The ordering kinetics in a tricritical system is studied for a model equation suitable for binary alloy such as Fe-Al mixture. The model consists of a coupled set of equations for the local order parameter field  $S(\underline{r}, t)$  and the local concentration  $X(\underline{r}, t)$  and reads

$$\partial_t S(\underline{r}, t) = -M \frac{\delta F\{S, X\}}{\delta S}$$

$$\partial_t X(\underline{r}, t) = LV^2 \frac{\delta F\{S, X\}}{\delta X}$$

where

$$F\{S, X\} = \int d\underline{r} \left[ \frac{h}{2} (\nabla X)^2 + \frac{k}{2} (\nabla S)^2 + W(S, X) \right]$$

$$W(S, X) = \frac{a}{2} X^2 - \frac{b}{2} S^2 + \frac{g}{4} S^4 + \frac{f}{2} S^2 X^2 - \mu X$$

All the coefficients  $a, b, \dots$  are chosen to be positive. The potential  $W(S, X)$  exhibits triple minima for  $2bf > ag$  and for an appropriate choice of the chemical potential  $\mu$ .

We have solved the above set of equations numerically in two dimensions with a periodic boundary condition. The parameters and the initial values for  $X$  and  $S$  are chosen to simulate the following three quenches:

- (1) from the disordered state at high temperatures to the ordered state between the second order transition line and the coexistence line.
- (2) from the ordered state to the inside of the coexistence line.
- (3) from the disordered state to the inside of the coexistence line.

We have observed the time evolution of the domain growth and have performed the scaling analysis of the scattering functions for  $X$  and  $S$ . The details will be described at the conference.